

Working with Planaria

ward's
science+

Working with Planaria

Introduction

Planaria are free-living aquatic flatworms known as Turbellarians, of the phylum Platyhelminthes (Greek for “flatworm”). Turbellarians are also known as “triclads”, which refers to their triple gut with a single anterior and two posterior branches.

Other characteristics of planaria include pigmentation which can be gray-brown, brown, white, or black; simple nervous system; and head with “eyes”. The eyes have no lenses and so cannot form an image, but they are sensitive to light. (To get a sense of how Planaria “see”, close your eyes and face a bright light.) Note, however, that planaria are negatively phototaxic—they avoid light; they hide under rocks, leaves, and other debris in aquatic habitats.

Planaria are carnivorous, eating living or dead animal matter. When they eat, they use their long, muscular pharynx, which they evert to almost one-half their body length.

Reproduction

Planaria reproduce both sexually and asexually. There are two methods of asexual reproduction: fragmentation and spontaneous “dropping tails”. Fragmentation usually begins with a transverse constriction just behind the pharynx, which increases until the two parts separate and move away from each other. The head grows a new tail, and the tail grows a new head. In spontaneous “dropping tails”, planaria in very stagnant water will frequently “drop” their tails, and the tails regenerate to form complete animals. This condition usually results in stunted animals, or dwarfs, which remain in this state until water conditions improve.

Because planaria are hermaphroditic—each animal possesses complete male and female systems—they reproduce sexually by producing “summer” eggs and “winter” eggs. Summer eggs are thin-shelled and transparent; winter eggs are usually black and on stalks. While summer eggs hatch quickly, winter eggs take longer to hatch; they may even remain dormant throughout the winter, and they are capable of withstanding unfavorable environmental conditions.

Collection

Planaria can be found on the underside of rocks, leaves, and other objects in the shallow water of streams, ponds, and rivers, as well as in aquatic vegetation such as *Elodea* and filamentous algae. The most common species found in running water are *Dugesia dorotocephala*, *Cura foremani*, and *Phagocata velata*; in standing water *Dugesia tigrina* and *Phagocata vernalis* are common. *Procotyla fluviatilis* can be found in running water, standing water, and even brackish water.

To collect planaria, wash off the underside of objects from a stream or pond bottom into a container of water. Planaria can also be collected by using “bait”—a cube of raw beef liver tied with a string and left to drift near submerged rocks for 15 or 20 minutes; shake off any planaria that adhere to the liver into a container of water. You may also remove a large amount of aquatic vegetation and place it in covered containers of water. As the oxygen content decreases, the planaria will rise to the surface where they can be easily removed.

Care and Feeding


Keep planaria in a covered shallow enamel pan; covered dishes may also be used. (Most species will also thrive in an aquarium.) Use pond or spring water; salts in tap water make it toxic to planaria. If tap water is used, remove salts with a water conditioner such as Fluval® Aquaplus ([470308-824](#)). Change the water every day or at least every other day. While planaria can survive in standing water, the water should be aerated ([470308-842](#)). Keep water at approximately 21°C or lower. If necessary, store the container with planaria on the bottom shelf of a refrigerator. To help the planaria avoid light, add cover such as slate or broken pieces of flower pots to the container. Clean the container once a week.

Feed the planaria once a week, making sure to remove any food that has not been eaten within three hours after feeding. We recommend performing a water change after each feeding.

White Planaria, *Procotyla fluviatilis*, should only eat small living crustaceans such as *Daphnia magna* ([470176-562](#)) and Brine Shrimp ([470180-300](#)). Other species can either eat the crustaceans or several other options including beef liver (which can be stored frozen), hard boiled eggs yolk, earthworm ([470210-928](#)) fragments, and chopped mealworms ([470183-784](#)).


Note: An interesting result of heavy feeding is the spontaneous “tail dropping” described above. If no food is available, a healthy planaria can survive for up to three months without harmful effects.

Figure A: *Dugesia tigrina*
diagrammatic representation




1. Ocellus
2. Auricle
3. Anterior branch of intestine
4. Gut diverticula
5. Areas of dispersed and concentrated pigment
6. Mouth
7. Everted pharynx (proboscis)
8. Posterior branch (2) of intestine

Figure B: Planarian
Section through pharynx



1. Dorsal epidermis with many rhabdites
2. Pharynx (proboscis)
3. Pharyngeal pouch
4. Posterior branch of intestine
5. Mouth, obliquely cut
6. Nerve Cord
7. Ciliated ventral epidermis
8. Diverticula of intestine
9. Parenchyma
10. Longitudinal muscle fibers
11. Circular muscle layer

Figure C: Planarian
Cross-section greatly magnified diagrammatic



1. Lumen of pharynx
2. Cilia of epithelial cells lining lumen
3. Longitudinal muscle fibers (longitudinal section)
4. Longitudinal muscle fibers (cross section)
5. Glandular cells
6. Circular muscle fibers
7. Pharyngeal pouch
8. Mouth, cut obliquely
9. Parenchyma (mesenchyme cells)
10. Rhabdite (rod-shaped structure present in epidermis of flatworms)
11. Ciliated ventral epidermis